SUPPLEMENTARY TABLE. Incidence rate* ratios of coccidioidomycosis, histoplasmosis, and blastomycosis, by selected patient characteristics — United States, 2019–2021

| | Coccidioidomycosis§ | | | Histoplasmosis [¶] | | | Blastomycosis** | | |
|--------------------------------|---------------------|-------------------|-------------------|-----------------------------|------------------|------------------|------------------|------------------|-------------------|
| Characteristic | 2019 | 2020 | 2021 | 2019 | 2020 | 2021 | 2019 | 2020 | 2021 |
| Sex | | | | | | | | | |
| Female | ref | ref | ref | ref | ref | ref | ref | ref | ref |
| Male | 1.1^{\dagger} | 1.2 [†] | 1.2 [†] | 1.3 [†] | 1.4 [†] | 1.4 [†] | 2.4 [†] | 1.8 [†] | 2.7 [†] |
| Age group (yrs) | | | | | | | | | |
| <5 | <0.1 [†] | <0.1 [†] | <0.1 [†] | 0.1 [†] | 0.1 [†] | 0.1 [†] | 0.1 [†] | 0.0 | <0.1 [†] |
| 5–19 | 0.3 [†] | 0.2 [†] | 0.2 | 0.4 [†] | 0.5 [†] | 0.5 [†] | 0.4 [†] | 0.3 [†] | 0.3 [†] |
| 20–39 | 0.7 | 0.6 [†] | 0.6 [†] | 0.7 | 0.8 [†] | 0.8 [†] | 1.0 | 0.6 | 0.6 [†] |
| 40–64 | ref | ref | ref | ref | ref | ref | ref | ref | ref |
| 65–79 | 1.4 [†] | 1.5 [†] | 1.5 [†] | 1.1 | 1.1 | 1.2 [†] | 1.1 | 1.2 | 1.1 |
| ≥80 | 1.3 [†] | 1.6 [†] | 1.5 [†] | 0.6 [†] | 0.6 [†] | 0.5 [†] | 0.5 | 0.6 | 0.6 |
| Race and ethnicity | | | | | | | | | |
| White, NH | ref | ref | ref | ref | ref | ref | ref | ref | ref |
| AI/AN, NH | 4.2 [†] | 4.4 [†] | 4.2 [†] | 0.9 | 1.2 | 1.5 | 6.4 [†] | 7.8 [†] | 7.4 [†] |
| Asian and NH/OPI, NH | 1.1 [†] | 1.0 | 0.9 [†] | 0.8 | 0.5 [†] | 0.5 [†] | 2.3 | 1.0 | 1.9 [†] |
| Black or African American, | 1.0 | 1.0 | 1.0 | 0.7 | 0.8 [†] | 0.8 [†] | 0.9 | 0.7 | 0.8 |
| NH | | | | | | | | | |
| Hispanic or Latino (all races) | 2.7^{\dagger} | 2.1 | 2.2 [†] | 0.9 | 0.6 [†] | 0.6 [†] | 1.3 | 1.0 | 1.3 |
| Other ^{††} | 1.1 | 1.5 [†] | 1.4 [†] | 1.5 [†] | 0.7 [†] | 0.8 | 0.3 | 0.1 | 0.1 |
| Total | NA | 0.9 [†] | 1.1 [†] | NA | 0.9 | 1.4 [†] | NA | 1.0 | 1.0 |

Abbreviations: AI/AN = American Indian or Alaska Native; NH = non-Hispanic; NH/OPI = Native Hawaiian or other Pacific Islander; ref = reference group; NA = not applicable.

^{*}Cases per 100,000 population. Summed state-specific denominators from 2019–2021 U.S. Census Bureau data were used to calculate incidence for the total grouping for each year.

[†]Statistically significant rate ratios based on a *p*-value of <0.05.

[§]Sex was missing for 68 cases, age was missing for 65 cases, and race and ethnicity were missing for 20,749 cases.

[¶]Sex was missing for five cases, age was missing for five cases, race and ethnicity were missing for 497 cases.

^{**}Race and ethnicity were missing for 48 cases.

^{††}Calculated on basis of two or more reported race categories.